

Agriculture & Food Security

Establishing symbiotic nitrogen fixation in cereals and other non-legume crops: The Greener Nitrogen Revolution

- David Dent and
- Edward Cocking

Agriculture & Food Security 2017:7 © The Author(s) 2017

Received: 19 August 2016 | **Accepted:** 14 December 2016 | **Published:** 1 March 2017

Abstract

Haber's invention of the synthesis of ammonia from its elements is one of the cornerstones of modern civilization. For nearly a century, agriculture has come to rely on synthetic nitrogen fertilizers produced from ammonia. This large-scale production is now supporting nearly half of the world's population through increased food production. But whilst the use of synthetic nitrogen fertilizers brought enormous benefits, including those of the Green Revolution, the world needs to disengage from our ever-increasing reliance on nitrogen fertilizers produced from fossil fuels. Their pollution of the atmosphere and water systems has become a major global environmental and economic concern. Naturally, legume crops such as peas and beans can fix nitrogen symbiotically by interacting with soil nitrogen-fixing rhizobia, bacteria that become established intracellularly within root nodules. Ever since this was first demonstrated in 1888, consistent attempts have been made to extend the symbiotic interaction of legumes with nitrogen-fixing bacteria to non-legume crops, particularly cereals. In 1988, a fresh impetus arose from the discovery of *Gluconacetobacter diazotrophicus* (*Gd*), a non-nodulating, non-rhizobial, nitrogen-fixing bacterium isolated from the intercellular juice of sugarcane. Subsequently, strains of *Gd* inoculated under specific conditions were shown to intracellularly colonize the roots and shoots of the cereals: wheat, maize (corn) and rice, as well as crops as diverse as potato, tea, oilseed rape, grass and tomato. An extensive field trials programme using a seed inoculum technology based on *Gd* (NFix[®]) indicates that NFix[®] is able to significantly improve yields of wheat, maize, oilseed rape and grasses, in both the presence and absence of synthetic nitrogen fertilizers. Evidence suggests that these benefits are accruing through a possible combination of intracellular symbiotic nitrogen fixation, enhanced rates of photosynthesis and the presence of additional plant growth factors. Here, we discuss the research events that have led to this important development and present results demonstrating the efficacy of NFix[®] technology in non-legume crops, in particular cereals.

Keywords

Gluconacetobacter diazotrophicus Nitrogen fertilizer Nitrogen fixation Intracellular colonization Yield benefits

A brief overview of nitrogen fertilizer production and pollution

The Haber–Bosch process for industrial ammonia production is considered by some to be the most important invention of the twentieth century. This important industrial development is partly responsible for the world's population explosion from 1.6 billion in 1900 to 6 billion in 2000 [1]. It has also been claimed that the use of 500 million tonnes of N fertilizer produced from ammonia each year, utilizing 1% of the world's energy and 3–5% of gas production [2], sustains around 40% of the global population [3]. However, the environmental costs of using ammonia-based nitrogen fertilizer across the globe are no longer considered sustainable. Nutrient excesses are particularly large in China, northern India, the

USA and Western Europe [4]. In the UK, agricultural nitrogen fertilizer use accounts for 66% of nitrous oxide emissions [5], whilst in Europe 5% of the population are exposed to drinking water contaminated with unsafe levels of nitrate [6].

In North America, agriculture is responsible for half of the nitrogen loading in Canadian surface waters [7, 8, 9], 75% of nitrous oxide emissions in the USA and for more than 1.5 million Americans drinking well water contaminated with nitrate [10]. Two-thirds of the coastal systems of the USA are moderately to severely impaired due to nutrient loading, with around 300 low-oxygen zones along the USA coastline. For example, the Mississippi River dumps 1.5 million metric tonnes of nitrogen into the Gulf of Mexico every year.

Addressing the pollution problems and damage caused by nitrogen fertilizer use is currently expensive making it unsustainable. Removing nitrates from water in Europe costs taxpayers between £130 and 650 per year. The economics of fertilizer use in European agriculture is questionable, given that the overall nitrous oxide damage costs £60–80 billion per year, a sum more than double the extra income gained from using nitrogen fertilizers in European agriculture [11].

The need for a sustainable alternative

There has been little argument that there is a need for more sustainable solutions to nitrogen fertilizer use. However, there is a shortage of genuine technical solutions available to address this global problem; hence, it remains difficult for governments to tackle the nitrogen issue.

There has been much support for developing technical solutions, not the least by the late Norman E. Borlaug, The Father of the Green Revolution. Hopes of biologically fixing nitrogen in cereals have been raised since the 1970s [12, 13], although these have yet to even be partially realized. Hence, an understandable reticence and caution has enveloped the whole field such that new developments are viewed with great scepticism, and even a reluctance to respect and accept results reflecting common standards of proof for the discipline. Such disparities in both interpretation and approach are indicative of an emerging paradigm shift [14, 15, 16].

Background to symbiotic nitrogen fixation in cereals

The introduction of symbiotic biological nitrogen fixation into cereals and other major non-legume crops would be regarded as one of the most significant contributions that biotechnology could make to agriculture. However, this has been recognized for many years as a major research challenge [17]. Currently, there are two strategic approaches used in attempts to achieve this long-standing aspiration. One is a long-term synthetic biology GM approach, engineering a nitrogen-fixing symbiosis from existing signalling and developmental mechanisms, to provide a suitable environment for rhizobial nitrogenase activity in the plant nodule [18]. The other, much shorter-term and simpler approach builds on the discovery that a non-rhizobial, naturally occurring nitrogen-fixing bacterium that fixes nitrogen in sugarcane is able to intracellularly colonize the root systems of cereals and other major crops [19]. In this approach, which is now at a field trial evaluation stage (Dent and Cocking in preparation), an adequate level of bacterial intracellular colonization and nitrogen fixation can be established throughout the plant without any need for nodulation. In such symbiotic nitrogen fixation, nitrogen-fixing bacteria establish an intracellular symbiosis with plants in which they fix nitrogen inside the cells of their host utilizing energy supplied by plant photosynthesis.

The need for intracellular colonization

Nodulated legumes in endosymbiosis with rhizobia are amongst the most important nitrogen-fixing systems in agriculture. In nodular cells, intracellular rhizobia, surrounded by a membrane derived by endocytosis from the plant plasma membrane, symbiotically fix nitrogen in cytoplasmic vesicles (symbiosomes). It is only by such intracellular colonization of nitrogen-fixing bacteria in these membrane-bound compartments (a state in which the bacteria are referred to as bacteroids) in the cytoplasm, that the prerequisites for symbiotic nitrogen fixation can be fulfilled. In symbiosomes, there is a reliable supply of metabolic substrates from plant biosynthesis which provides (a) sufficient energy and reducing conditions, (b) protection against elevated oxygen concentrations that inhibit nitrogen fixation, (c) a pathway for the transport of nitrogen fixation products to the plant host, (d) development of membrane systems for bidirectional transport between the host and bacteria and endosymbiont and (e) protection against competitive or antagonistic bacteria in the environment [20].

However, the actual formation of nodules is not an essential requirement for the establishment of intracellular plant endosymbiotic nitrogen fixation. In non-legume angiosperm *Gunnera* plants, the nitrogen-fixing cyanobacterium, *Nostoc*, enters mucilage-secreting glands found on the stem in the axis of the leaves. It invades the gland cells and becomes intracellular in membrane-bound vesicles in the cytoplasm, similar to symbiosomes in rhizobium–legume symbiosis but without nodule formation [21, 22]. The cyanobacteria divide within the intracellular vesicles and differentiate into heterocysts. These heterocysts possess a double-layered envelope of polysaccharide and glycolipids which act as a barrier to oxygen diffusion. The quantity of nitrogen fixed by the intracellular heterocyst symbiont and transferred to the *Gunnera* host plant is sufficient to satisfy its nitrogen requirements [23].

Bacteria which can colonize plants intracellularly have now been identified in a wide range of plant species including non-nodulating legumes colonized by rhizobia [24], peach palm [25], scotch pine [26], Engelmann spruce and limber pine [27], banana [28, 29] and cactus, four of which include nitrogen-fixing bacteria [27, 30].

The ability of nitrogen-fixing bacteria, other than *Nostoc*, to endosymbiotically fix nitrogen in symbiosomes in non-legumes, without the need for nodulation is of major evolutionary interest. Also the ability of rhizobia to fix nitrogen in 13 species of non-nodulating legumes (species of the subfamilies Caesalpinioideae, Papilionoideae and Mimosoideae [31, 32]) has important implications from an evolutionary perspective [32]. There is sufficient evidence available reporting that nodule formation is not a necessary prerequisite for intracellular plant nitrogen fixation, even in the rhizobia–legume symbiosis. The Leguminosae nodulation seems to occur amongst the more specialized subfamilies. It is important to note that the host plant itself encodes the genetic developmental programme responsible for the development of the nodule tissues and for regulation of the process, with organogenesis being triggered by the rhizobial micro-symbiont [33]. This raises questions in regard to the means of cell entry in non-nodular and non-leguminous symbioses.

Entry of nitrogen-fixing bacteria

Until now, it has been unclear as to how nitrogen-fixing bacteria are able to breach the cell walls of plants to establish endosymbiosis; more is known about the uptake of rhizobia. They are endocytosed into plant cells in which they become surrounded by a peribacteroid membrane to create the symbiosome structure referred to above. Endocytosis is only possible after bacteria have penetrated the cell wall and are able to interact with the plant cell plasma membrane. Interestingly, forms resembling bacteroids have long been reported inside the roots of non-nodulating *Gleditsia* species [34, 35, 36]. This finding is supported by work using scanning electron microscopy in *G. triacanthos* and other species [31]. Penetration of these species (*G. triacanthos*, *Cassia fistula*, *C. grandis* and *Senna tora*) cell wall is most likely to occur through the formation of infection threads in the root hairs [31, 37], whereas other mechanisms of entry are also evident with rhizobia. Particularly pertinent in this regard is the

production and isolation of cell-bound cellulases of white clover rhizobia that can erode a complete hole, slightly larger than the bacterial cell, that traverses the non-crystalline wall region of white clover root hair apices [38]. Alternatively, plant cell wall-degrading enzymes induced in response to rhizobia may be involved.

Further insights into the entry of nitrogen-fixing bacteria into plant cells and the establishment of intracellular colonization have been obtained from an extensive range of experiments extending between 1960 and the mid-1990s, using enzymatically isolated plant protoplasts. The isolation of plant protoplasts using cellulases to digest the cell wall opened a whole new range of studies in plant cellular and molecular biology. What made the isolated protoplast so powerful as an experimental system was that the enzymatic removal of the cell wall left the cell surface membrane (the plasma membrane) fully exposed as the only barrier between the external environment and the interior of the cell. The accessibility of the plasma membrane meant that experiments could be designed to investigate and manipulate the properties of this membrane, in a way that was not possible with walled cells [39]. It was therefore possible to demonstrate the endocytotic uptake of rhizobia into pea leaf protoplasts [40]. This seminal finding showed clearly the need to select bacteria for symbiotic interactions that are both nitrogen fixing and cellulase producing. The origin of the membrane-bound vesicles containing rhizobia was directly analogous to that of vesicles enclosing rhizobia, which enter root nodule cells via the normal infection thread. This highlighted the importance of cell wall degradation, without loss of cell viability, to establish the cell as a habitat for nitrogen-fixing bacteria in membrane-bound vesicles destined to become symbiosomes.

The ability to isolate nodule protoplasts containing symbiosomes with bacteroids [41], coupled with the ability to fuse these protoplasts with protoplasts isolated from non-legumes, provided another way of establishing rhizobial symbiosomes in non-legume crops for the production of nitrogen-fixing hybrid cells, possibly leading to the formation of nitrogen-fixing somatic hybrid plants. A wide range of heterokaryons between legumes and non-legumes was produced [42]. However, it proved impossible to maintain the ongoing viability and cell division of these heterokaryons.

Increasing the host range of rhizobia

Establishing intracellular colonization and symbiosome formation in non-legume crops could result from increasing the host range of rhizobia. In the soil–root surface, the rhizosphere, developing root hairs play an important role in symbiotic recognition. Rhizobia, and especially the Nod factors (signal lipo-chitooligosaccharides) they secrete, stimulate the re-orientation of root hair cell wall growth, resulting in curled root hairs. Nod factors promote the formation of infection threads, and it is through these tubular structures that rhizobia enter most legumes. For temperate legumes such as clover, there is a very tight control of *Rhizobium* specificity. A barrier to this specificity could be removed by enzymatically degrading the cell wall at the apices of root hairs in white clover, using a cellulase–pectolyase enzyme mixture. Subsequent inoculation with rhizobia in the presence of polyethylene glycol enabled rhizobia to induce the formation of nitrogen-fixing nodules on white clover (a non-host). This cellulase–pectolyase enzyme treatment is also known to release protoplasts from the tips of root hairs of legume and non-legume crop plants and provides an opportunity for rhizobial endocytosis via root tip interaction, thus identifying the apical cell wall of root hairs as a target site for the nodulation of cereals and other non-legume crops [43].

In rice, it was shown that rhizobia could induce formation of nodule structures provided the rice seedling root hairs were treated with cell wall-degrading enzymes and polyethylene glycol. These nodule structures consisted of several cell layers enclosing scattered centrally located cells, with rhizobia mainly located in spaces between the cell layers, and in dead cells of the central region, there was negligible nitrogenase activity [44]. Similar enzyme treatment and inoculation of oilseed rape also produced nodular structures morphologically resembling clover nodules. However, only dead cells were invaded by rhizobia and there was only a very low nitrogenase activity [45]. From these attempts, it became progressively evident in the 1990s that it was not possible, using these novel inoculation

procedures to imitate in non-legume crops the organized nodule development pattern, characteristic of legume nodulation by rhizobia. Nor was it possible to obtain the membrane-bound rhizobia colonization of the living nodular cells required for endosymbiotic nitrogen fixation.

This failure to achieve effective nodulation of non-legume crops by rhizobia led to a search for a non-nodular niche for endophytic nitrogen fixation. This was comparable to the situation arising from the naturally occurring non-nodular colonization of intercellular spaces and xylem of sugarcane by endophytic nitrogen-fixing bacteria. Some sugarcane varieties are able to obtain over 60% of their nitrogen from plant associated biological nitrogen fixation [46].

The use of other nitrogen-fixing bacteria

Rhizobia are now regarded as being inadequate to establish symbiotic nitrogen fixation in non-legume crops, because they do not colonize living cells and their nitrogen-fixing capability is inhibited by oxygen [47, 48]. Regulating the flux of oxygen to endosymbiotic rhizobia has been suggested to be an almost insuperable challenge to the dream of introducing rhizobial nitrogen-fixing symbiosis effectively into non-legume crops [48]. The use of other nitrogen-fixing bacteria such as *Azoarcus* spp., an endophyte of Kallar grass, and *Klebsiella* spp., an endophyte of maize, highlighted the readiness with which endophyte colonization of intercellular spaces and dead xylem cells can occur following their inoculation of non-legume crops. However, like with rhizobia in non-legumes, there was no evidence for any intracellular colonization of living cells in any of the plants investigated and therefore no opportunity for the establishment of symbiotic nitrogen fixation [49].

Nitrogen fixation in sugarcane

The long-term continuous cultivation of sugarcane in Brazil uses low N fertilizer inputs without apparent depletion of soil N response. This led to the suggestion that nitrogen-fixing bacteria associated with the sugarcane might be the source of the agronomically significant nitrogen inputs to this crop [50]. One of these was a new species of nitrogen-fixing non-rhizobial, aerobic bacterium that had been isolated in 1988 from Brazilian sugarcane roots and stems [51]. The same species of bacteria was later extracted in large numbers from the xylem of Mexican sugarcane cultivars, which are also colonized intercellularly [52]. This bacterium, later renamed *Gluconacetobacter diazotrophicus* (*Gd*), has many characteristics that made it attractive for the establishment of intracellular symbiotic nitrogen fixation, without the need for nodulation, paralleling the situation in the angiosperm *Gunnera* with *Nostoc* [21]. *Gd* is known to produce cellulases, hemicellulases (xyloglucanases) and pectinases, which could facilitate the penetration of plant cell walls [53], all of which are formed in the presence of sucrose which is involved in the establishment of the *Gunnera*–*Nostoc* endosymbiosis [54].

Gluconacetobacter diazotrophicus (*Gd*), ‘this most extraordinary diazotroph’ [55], is a small, 0.7–0.9 by 1–2 μm , motile, gram-negative aerobic rod showing pellicle formation. Best growth occurs with high sucrose or glucose concentrations (10% w/v) [56]. It secretes an enzyme, levansucrase, which hydrolyses sucrose and catalyses the formation of the exopolysaccharide fructan levan [57]¹ which limits oxygen diffusion. This protects nitrogenase activity from excessive oxygen flux, enabling the bacteria to fix nitrogen even if the pO_2 is not much lower than tropospheric levels [58]. Detoxification of reactive oxygen species may also occur [59]. *Gd* possesses no nitrate reductase, and nitrogen fixation is not inhibited by high concentrations of nitrate (25 mM). The nitrogenase activity of *Gd* is only partially inhibited by ammonium, implying that *Gd* can continue fixing nitrogen inside the plant even when nitrate (and ammonium)-containing fertilizers are applied. Another unique feature of this bacterium is its ability to excrete almost half of the fixed nitrogen as ammonium, which becomes potentially available to plants [60].

The intracellular establishment of *Gd*

The primary cell walls of plant meristems consist of β -1,4-glucan-linked chains of cellulose that coalesce via hydrogen bonding to form polysaccharide microfibrils [61]. The action of *Gd* cell wall-degrading enzymes might weaken the primary cell walls of root meristems sufficiently, enabling *Gd* to interact directly with the plasma membrane of root cell protoplasts. Endocytotic uptake of *Gd* into membrane-bound vesicles in the cytoplasm could then result in the intracellular establishment of *Gd* in symbiosomes, similar to rhizobia in root nodule cells [19]. Using seedlings of maize, rice, wheat, oilseed rape, tomato, white clover and *Arabidopsis* grown aseptically in sucrose-containing culture media, inoculation with very low numbers of *Gd* has resulted in extensive intracellular colonization of root meristems within membrane-bound vesicles [19]. *Gd* subsequently spreads intracellularly from the root tip meristem into cells of the elongating region of the root and into leaf cells of the shoot system in which *Gd* associates closely with chloroplasts. This raises the possibility that energy might be supplied directly from chloroplasts to nitrogen-fixing endosymbiotic bacteria within the same cell.

The generality of the intracellular colonization of this wide range of plant species by *Gd*, contrasting with the high degree of rhizobial strain specificity for most legume nodule interactions, may be due to the very close similarity in primary wall composition of root meristems throughout the plant kingdom [19]. This discovery of plant systemic intracellular colonization by *Gd* sets the stage for investigating their possible utilization to reduce synthetic nitrogen fertilizer application used in agriculture, and any additional effects on intrinsic yield potential.

Implications for evolution

The ability of *Gd* to systematically colonize both non-legumes and legumes intracellularly and without nodulation could provide new insights into the evolutionary origin of symbiotic nitrogen fixation in plants. Postgate [62] has posed the question as to why there is no such thing as a nitrogen-fixing plant and why do plants have to form a symbiotic association with nitrogen-fixing bacteria to gain access to atmospheric nitrogen? In terms of energetics, nitrogen fixation should not present an evolutionary obstacle to plants. Eight molecules of ATP are required for every 1/2 molecule of N_2 reduced to NH_3 by the bacterial nitrogenase. Additional six ATP molecules are required as a reductant, resulting in a need of 14 ATP molecules per NH_3 produced. Plants usually assimilate their nitrogen by reduction of nitrate that requires 12 molecules of ATP to provide one molecule of NH_3 . Thus, nitrogen fixation is only marginally more demanding than nitrate reduction in terms of energy consumption.

It is surprising that no plant has yet become independent of bacteria in relation to N uptake [62]. The development of a 'diazoplast', a new organelle analogous to a chloroplast, would help plants overcome the genetic and physiological bacterial-dependent nitrogen fixation problems. The diazoplast could be acquired similarly to a chloroplast by an accretion of an endosymbiotic prokaryote into the plants genome. The inoculation of plants with *Gd*, in the presence of sucrose, results in its establishment intracellularly in root meristems and then systemically in roots and shoots within plant membrane-bound symbiosome-like vesicular compartments in the cytoplasm. These structures are likely the location for the formation of diazoplasts and the emergence of autonomous nitrogen-fixing plants by 'giving evolution a push in a direction in which it is already poised to go' [62]. It would now seem likely that the independent evolution of intracellular non-nodular nitrogen fixation, thought at present to be restricted to the Gunneraceae [63], is not unique but that such non-nodular nitrogen fixation can be extended to a very wide range of plant families [25, 26, 27, 32]. Interestingly the phylotype of some conifers has been found to be related to *Gd* and other nitrogen-fixing acetic acid bacterial endophytes including *Acetobacter*, *Acidomonas*, *Kozakia* and *Asaia* [27]. This suggests that non-nodular, endocytotic, intracellular meristem cell colonization by a nitrogen-fixing and cellulase producing bacterium, such as *Gd*, is the actual single evolutionary innovation that drove the deep evolution of symbiotic nitrogen fixation in plants [64].

Agricultural value of *G. diazotrophicus*

The criteria for evaluating the practicality of using any nitrogen-fixing bacterium in agriculture need to at least match those applied for the assessment of the efficacy of rhizobia/legume associations. Leaving aside the need for colonization, intracellularity and the development of symbiosomes (considered above), BNF studies to demonstrate nitrogen fixation should look for increases relative to controls associated with the following criteria (expanded on below):

- The measurement of foliage greenness/chlorophyll content
- The measurement of the percentage N in the biomass
- Labelled N studies
- *Nif* minus mutants
- The measurement of nitrogenase activity by e.g. ARA
- Demonstration of field efficacy and yield benefits in nitrogen poor soils in the absence and/or presence of a nitrogen fertilizer

Foliage greenness/improved chlorophyll content: There is a very close link between chlorophyll and nitrogen content [65, 66, 67]. Studies with legume/rhizobia associations have reported enhanced leaf chlorophyll content in chickpea [68], cowpea [69] and common bean [70] when the associations were established. Similarly, chlorophyll levels have been demonstrated to increase in sorghum colonized by *Gd* [71] and leaf greenness increased in *Lolium perenne* with *Gd* IMI501986 relative to uninoculated controls (Patent Number: WO2016/016629).

Percentage N in leaves: A 40% increase in leaf N in sorghum has been demonstrated when co-inoculated with AM fungi [71]. Trials by Azotic Technologies have shown a 43% increase in leaf N of inoculated *L. perenne* relative to uninoculated field plots (Dent and Cocking in preparation).

Nitrogenase activity demonstrated by ARA: The acetylene reduction assay (ARA) has been used for many years to demonstrate nitrogen fixation by rhizobia in root nodules. For *Gd*, similar assays have successfully focused on nitrogen fixation capability of bacteria in culture [72, 73], but when attempted for inoculated maize plants there was no evidence of nitrogen fixation [74].

Nitrogen fixation demonstrated by *Nif* minus studies: The ability to use *Nif* mutants in a bacterium and therefore the loss of nitrogen-fixing ability is the key indicator of nitrogen fixation *in planta*. For *Gd*, this was achieved by demonstrating that the shoot nitrogen content of sugarcane plants inoculated with wild-type Pal5 was significantly higher than in uninoculated control plants, or plants inoculated with the *Nif*-mutant Mad3A in nitrogen-limiting conditions [75].

Nitrogen fixation demonstrated by labelled nitrogen ($^{15}\text{N}_2$) studies: A $^{15}\text{N}_2$ incorporation experiment demonstrated that *Gd* Pal5 wild-type strains actively fixed nitrogen inside sugarcane plants, whereas *Nif* mutants did not [75]. Another $^{15}\text{N}_2$ incorporation experiment also demonstrated nitrogen fixation by *Gd* in sugarcane [73].

Plant growth and yield benefits: Plant growth and yield parameters (seed yield, plant size, flowering times) have been shown to be markedly improved in rhizobia inoculated versus uninoculated legume crops for soybean [76, 77], pea and lentils [78], common bean [79], chickpea [80], mung bean [81] and cowpea [69]. Similarly with *Gd*, tomato plants inoculated with the Pal5 strain significantly increased both number and weight of tomato fruit production as compared to non-inoculated controls [82]. Biomass accumulation following inoculation with 14 strains of *Gluconacetobacter* spp. was well matched with corresponding nitrogen fixation in all tissues of sugar beet [83]. Inoculation with two strains of *Gd* (Pal5, UAP5541), which colonized the roots and shoots of sorghum and wheat, increased both the shoot and root dry weight of sorghum, whereas they had no effect on either shoot or root dry weight of maize [84, 85]. Interestingly, Pal5 inoculation was also unable to relieve the N-deficiency symptoms of unfertilized maize in either the field or the greenhouse [86]. However, recent independent

field trial research utilizing proprietary NFix[®] technology (Patent Number: WO2016/016629) based on *Gd* demonstrated significant yield improvements of around 1 tonne per hectare in both maize and wheat (Figs. 1, 2, for raw data see Additional file 1).




Fig. 1

Yield of maize inoculated with NFix[®] against uninoculated controls in an independent N fertilizer trial. This N fertilizer trial was conducted by independent third-party contract researchers on maize variety PR39F58 in Germany. The untreated seed control was compared with the NFix[®] treatment at five different levels of N fertilizer application. *Bars on each point* represent standard errors of the means of yields from six plots (60 m²) in a split-plot design (main plot: seed treatment; subplot: nitrogen fertilizer). The *dotted line* indicates when the yield of the full recommended rate of N fertilizer* according to the total available N in the soil pre-sowing is equivalent to the yield of the NFix[®] treatment




Fig. 2

Yield of wheat inoculated with NFix[®] against uninoculated controls in an independent N fertilizer trial. This N fertilizer trial was conducted by independent third-party contract researchers in the UK on wheat variety Mulika. The untreated seed control was compared with the NFix[®] treatment at five different levels of N fertilizer application. *Bars on each point* represent standard errors of the means of yields from four plots (24 m²) in a split-plot design (main plot: seed treatment; subplot: nitrogen fertilizer). The *dotted line* indicates when the yield of the full recommended rate of N fertilizer* according to the total available N in the soil pre-sowing is equivalent to the yield of the NFix[®] treatment

Photosynthetic rates: The rhizobia–legume symbiotic association is known to contribute benefits to the plant greater than just nitrogen fixation. Indeed, even with the cost to the plant of reduced photosynthate by as much as 14% [87], rhizobia have been demonstrated to increase the rate of photosynthesis and other associated measures in common bean (intercellular CO₂ concentration and the transpiration rate) [70], peas [88] and soybean [87]. Rhizobial colonization of the non-legume crop rice increased photosynthetic rate, stomatal conductance, transpiration velocity, water utilization efficiency and flag

leaf area. Furthermore, it was shown that higher levels of the growth hormone indole-3-acetic acid (IAA) and gibberellic acid (GA) were accumulated [89]. The only research that has been conducted in this field with *Gd* has involved the inoculation of *Arabidopsis thaliana*. In this study, *Gd* had beneficial effects on photosynthesis [90]. Whole-canopy photosynthesis levels were 68% higher in inoculated plants compared to uninoculated plants. However, transpiration rates were 15.5% lower. As a consequence of greater carbon uptake and lower water loss through transpiration, inoculated plants showed increases of 94.5% in whole-canopy water-use efficiency compared to non-inoculated plants. These results suggest that *Gd* positively affects plant physiology by increasing the photosynthetic capacity and improving water-use efficiency [90], which goes some way to explaining the parallel yield relationships across all levels of N fertilizers, shown in Figs. 1 and 2.

Conclusions

The need for an improved means of delivering nitrogen to cereals and other non-legume crops is crucial for the future of sustainable agriculture, including the reduction of ammonia, nitrate and nitrous oxide pollution but ensuring food security [91].² The promise of biological nitrogen fixation for cereals proffered in the 1980s may now be realized, however, not as originally envisaged through rhizobial association with genetically manipulated root nodules on wheat for instance, but rather through appropriate application of the naturally occurring nitrogen-fixing endophyte *Gd* to the plant using products, such as NFix[®].

The necessary ability for intracellular colonization was established initially in 2006 with *Gd* in a range of crop species [19]. This has now been independently validated with a number of nitrogen-fixing bacteria for a range of plants (e.g. [27]), and the enzymes required for cell entry determined. Measures used to establish nitrogen fixation and plant growth benefits in rhizobia–legume symbiosis have been successfully applied to *Gd*, similarly demonstrating nitrogen fixation and significant crop yield benefits, the latter even in the presence of full recommended rates of nitrogen fertilizer. The development of a nitrogen-fixing endophytic bacterium that can be applied to all staple food crops and substitute for mineral nitrogen fertilizers, whilst delivering yield benefits, is a significant major development and heralds the prospect of a *Greener Nitrogen Revolution!*

Footnotes

1

With the recent discovery of nitrogen fixation in sugarcane, the development of nitrogen fixation in other grasses now seems within our grasp in the near future. This development will likely not be based on the introduction of *nif* genes into grasses or on the induction of nodules on grass roots that harbour diazotrophs. This development will most likely be based on the successful inoculation of diazotrophs that infect and inhabit the roots and stems of grass species in a manner similar to the sugarcane system.

2

Creating new strains of rice, wheat and corn that fix their own nitrogen could achieve in the twenty-first century is what the Haber–Bosch breakthrough managed for the twentieth and without the serious environmental drawbacks of industrial ammonia production. Environmentalists should not be scared of this prospect; they should welcome it. There can be no more important task than feeding people whilst protecting the planet. We must use the best of science and technology to help us to achieve this vital aim.

Abbreviations

Gd: *Gluconacetobacter diazotrophicus*
GM: genetically modified
N: nitrogen
ATP: adenosine triphosphate
ARA: acetylene reduction assay
BNF: biological nitrogen fixation
AM fungi: arbuscular Mycorrhizal fungi
IAA: indole-3-acetic acid
GA: gibberellic acid

Declarations

Authors' contributions

DD and EC wrote the manuscript. Both authors have read and approved the final manuscript.

Acknowledgements

Thanks to Dr. Pedro Carvalho for the analysis and presentation of Figs. [1](#) and [2](#) and the Azotic Technologies Ltd. and Koppert BV R&D teams for their inputs to this paper, particularly Dr. Katrin Schwarz and Daniel Rooney for their editorial assistance.

Dedicated to the pioneering research and memory of José de Jesús Caballero-Mellado (1953–2010).

Competing interests

DD is a Founder Director of Azotic Technologies Ltd. and Chief Technology Officer and hence has been responsible for the development of the NFix[®] formulation cited in the text and the patent WO2016/016629.

Availability of data and materials

All data generated or analysed during this study are included in this published article [and its supplementary information files].

Funding

This work was funded by Azotic Technologies Ltd.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

Additional files

[40066_2016_84_MOESM1_ESM.xlsx](#) Additional file 1.

References

1. Smil V. Detonator of the population explosion. *Nature*. 1999;400:415. [View ArticleGoogle Scholar](#)
2. Smith BE. Structure. Nitrogenase reveals its inner secrets. *Science*. 2002;297(5587):1654–5. [PubMedView ArticleGoogle Scholar](#)
3. Fryzuk MD. Ammonia transformed. *Nature*. 2004;427(6974):498–9. [PubMedView ArticleGoogle Scholar](#)
4. Foley JA, Ramankutty N, Brauman KA, Cassidy ES, Gerber JS, Johnston M, et al. Solutions for a cultivated planet. *Nature*. 2011;478:337–42. [PubMedView ArticleGoogle Scholar](#)
5. DEFRA. Guidance: Air pollution from farming: preventing and minimizing. Department for Environment, Food and Rural Affairs and Environment Agency. <https://www.gov.uk/guidance/reducing-air-pollution-on-farms> (2013). Accessed 28 Apr 2016.
6. van Grinsven H, Ward MH, Benjamin N, de Kok TMCM. Does the evidence about health risks associated with nitrate ingestion warrant an increase of the nitrate standard for drinking water? *Environ Health*. 2006;5:5–26. [View ArticleGoogle Scholar](#)
7. Chambers PA, Kent R, Chariton MN, Guy M, Gagnon C, Roberts E, et al. Nutrients and their impact on the Canadian environment. Agriculture and Agri-Food Canada, Environment Canada, Fisheries and Oceans Canada, Health Canada, and Natural Resources Canada, Ottawa, ON, Canada. <http://publications.gc.ca/collections/Collection/En21-205-2001E-2.pdf> (2001). Accessed 16 Apr 2016.
8. Janzen HH, Beauchemin KA, Bruinsma TW, Campbell CA, Desjardins RL, Ellert BH, et al. The fate of nitrogen in agroecosystems: an illustration using Canadian estimates. *Nutr Cycl Agroecosyst*. 2003;67:85–102. [View ArticleGoogle Scholar](#)
9. Rasouli S, Whalen JK, Madramootoo CA. Review: reducing residual soil nitrogen losses from agroecosystems for surface water protection in Quebec and Ontario, Canada: best management practices, policies and perspectives. *Can J Soil Sci*. 2014;94:109–27. [View ArticleGoogle Scholar](#)
10. Davidson EA, David MB, Galloway JN, Goodale CL, Haeuber R, Harrison JA, et al. Excess nitrogen in the US Environment: trends, risks and solutions. *ESA Issue Ecol*. 2012;15:1–16. [Google Scholar](#)
11. INRA Research. *The European Nitrogen Assessment: Sources, effects and policy perspectives*. In: Sutton MA, Howard CM, Erisman JW, Bleeker GBA, Grennfelt P, van Grinsven H, et al., editors. INRA Research April 2011. Cambridge: Cambridge University Press; 2011. p. 664. [Google Scholar](#)
12. Simmonds J. *Community matters: a history of biological nitrogen fixation and nodulation research 1965–1995*. Ph.D. Thesis Rensselaer Polytechnic Institute. Troy, New York: UMI Number 3299478. 2008.
13. Rogers C, Oldroyd GED. Synthetic biology approaches to engineering the nitrogen symbiosis in cereals. *J Exp Bot*. 2014;65(8):1939–46. [PubMedView ArticleGoogle Scholar](#)
14. Kuhn TS. *The essential tension: Tradition and Innovation in scientific research*. In: Kuhn TS, editor. *Essential tension*. Chicago: The University of Chicago Press; 1977. p. 225–9. [Google Scholar](#)
15. Dent DR. *Experimental Paradigms*. In: Dent DR, editor. *Integrated pest management*. London: Chapman and Hall London; 1995. p. 172–208. [Google Scholar](#)
16. Dent DR. *Panaceas, paradigms and pragmatism*. In: Dent DR, editor. *Insect pest management*. Wallingford: CABI; 2002. p. 327–30. [Google Scholar](#)
17. Conway G. *The Doubly Green Revolution: food for all in the 21st century*. London: Penguin Books; 1997. p. 335. [Google Scholar](#)
18. Oldroyd GED, Dixon R. Biotech solutions to the nitrogen problem. *Curr Opin Biotechnol*. 2014;26:19–24. [PubMedView ArticleGoogle Scholar](#)
19. Cocking EC, Stone PJ, Davey MR. Intracellular colonisation of roots of *Arabidopsis* and crop plants by *Gluconacetobacter diazotrophicus*. *Vitro Cell Dev Biol Plant*. 2006;42(1):74–82. [View ArticleGoogle Scholar](#)

20. Quispel A. Critical evaluation of prospects for nitrogen fixation with non-legumes. *Plant Soil*. 1991;137:1–11. [View ArticleGoogle Scholar](#)
21. Werner D. Symbiosis of plants and microbes. In: Tikhonovich IA, Provorov NA, Romanov VI, Newton WE (eds) *Nitrogen fixation: Fundamentals and applications: proceedings of the 10th international congress on nitrogen fixation*. St Petersburg: Springer; 1995. BV:241.
22. (Unknown surname). *The Gunnera-Nostoc Association: Why should you let me in?* <https://sciweek.wordpress.com/2012/09/12/> (2012). Accessed 20 May 2016.
23. Sylvester WB, Smith DR. Nitrogen fixation by *Gunnera Nostoc* Symbiosis. *Nature*. 1969;224:1231. [View ArticleGoogle Scholar](#)
24. Sambukumar S, Affeldt K, Friedrich D, Rausch S, Reddy P, James E, et al. (2015) Exploring the Rhizobial interactions with non-nodulating legume *Gleditsia triacanthos*. In: 23rd North American conference on symbiotic nitrogen fixation, 6–10 Dec 2015. Ixtapa, Mexico.
25. de Almeida CV, Andreote FD, Yara R, Tanaka FAO, Azevedo JL, de Almeida M. Bacteriosomes in axenic plants: endophytes as stable endosymbionts. *J Microbiol Biotechnol*. 2009;25:1757–64. [View ArticleGoogle Scholar](#)
26. Pirttilä AM, Laukkanen H, Pospiech H, Myllylä R, Hohtola A. Detection of intracellular bacteria in the buds of Scotch pine (*Pinus sylvestris*) by *in situ* hybridization. *Appl Environ Microbiol*. 2000;66:3073–7. [PubMedPubMed CentralView ArticleGoogle Scholar](#)
27. Carrel AA, Frank AC. *Pinus flexilis* and *Picea engelmannii* share a simple and consistent needle endophyte microbiota with a potential role in nitrogen fixation. *Front Microbiol*. 2014;5(333):1–11. [Google Scholar](#)
28. Thomas P, Reddy MK. Microscopic elucidation of abundant endophytic bacteria colonizing the cell wall—plasma membrane perispace in the shoot-tip tissue of banana. *AoB PLANTS*, 5:plt011. 2013. doi:[10.1093/aobpla/plt011](https://doi.org/10.1093/aobpla/plt011).
29. Thomas P, Sekhar AC. Live cell imaging reveals extensive intracellular cytoplasmic colonisation of banana by normally non-cultivable endophytic bacteria. *AoB PLANTS*, 6:plu002. 2014. doi.org/[10.1093/aobpla/plu002](https://doi.org/10.1093/aobpla/plu002).
30. White JK, Torres MF, Johnson H, Irizarry I, Chen Q, Zhang N, et al. Intracellular colonization and oxidative lysis of bacteria in vascular plant seedling tissues. *ResearchGate*. https://www.researchgate.net/publication/247778278_Intracellular_colonization_and_oxidative_lysis_of_bacteria_in_vascular_plant_seedling_tissues (2013). Accessed 22 Apr 2016.
31. Bryan JA. Nitrogen-fixing Leguminous trees and shrubs: a basic resource of agroforestry. In: Ashton MS, Montagnini F, editors. *The silvicultural basis for agroforestry systems*. Abingdon: CRC Press, Taylor & Francis Group; 1995. p. 41–60. [Google Scholar](#)
32. Bryan JA, Berlyn GP, Gordon JC. Towards a new concept of the evolution of symbiotic nitrogen fixation in the Leguminosae. *Plant Soil*. 1996;186:151–9. [View ArticleGoogle Scholar](#)
33. Trichene L, James EK, Sandal N, Stougaard J. Spontaneous root-nodule formation in the model legume *Lotus japonicus*: a novel class of mutants nodulates in the absence of Rhizobia. *Mol Plant Microbe Interact*. 2006;19(4):373–82. [View ArticleGoogle Scholar](#)
34. Feher D, Bolcor R. Untersuchungen über die bakterielle Wurzelsymbiose einiger Leguminosenhölzer. *Planta*. 1926;2:406–13. [View ArticleGoogle Scholar](#)
35. Friesner GM. Bacteria in the roots of *Gleditsia triacanthos*. *Proc Indiana Acad Sci*. 1926;34:215–24. [Google Scholar](#)
36. Rothschild D I de. Estudio preliminar sobre la presencia de bacterias en la raíz de *Gleditsia amorphoides* y *G. triacanthos*. *Comunicaciones del Museo Argentino de Ciencias Naturales “Bernardino Rivadavia” e Instituto Nacional de Investigacion de las Ciencias Naturales. Botanica*. 1967;2:9–14.
37. Allen ON, Allen EK. *The Leguminosae: a source book of characteristics, uses and nodulation*. Madison: University of Wisconsin Press/London: W/Macmillan Publishing; 1981.
38. Mateos PF, Baker DL, Petersen M, Velázquez E, Jiménez-Zurdo JI, Martínez-Molina E, et al. Erosion of root epidermal cell walls by *Rhizobium* polysaccharide-degrading enzymes as related to primary host infection in the *Rhizobium* legume symbiosis. *Can J Microbiol*. 2001;47(6):475–87. [PubMedGoogle Scholar](#)

39. Cocking EC. The contribution of the study of plant protoplasts to cellular and molecular biology. *J Plant Res.* 1983;3:177–80. [Google Scholar](#)
40. Davey MR, Cocking EC. Uptake of bacteria by isolated higher plant protoplasts. *Nature.* 1972;239:455–6. [View ArticleGoogle Scholar](#)
41. Davey MR, Cocking EC, Bush E. Isolation of legume root nodule protoplasts. *Nature.* 1973;244:460–1. [View ArticleGoogle Scholar](#)
42. Davey MR, Cocking EC. Tissue and cell culture of bacterial nitrogen fixation. In: Rao S, editor. *Recent advances in biological nitrogen fixation.* New Delhi, Bombay, Calcutta: Oxford and IBH Publishing Co; 1980. p. 281–324. [Google Scholar](#)
43. Al-Mallah MK, Davey MR, Cocking EC. Enzymatic treatment of clover root hairs removes a barrier to rhizobium-host specificity. *Nat Biotechnol.* 1987;5:1319–22. doi: [10.1038/nbt1287-1319](https://doi.org/10.1038/nbt1287-1319). [View ArticleGoogle Scholar](#)
44. Al-Mallah MK, Davey MR, Cocking EC. Formation of nodular structures on rice seedlings by *Rhizobia*. *J Exp Bot.* 1987;40:473–8. [View ArticleGoogle Scholar](#)
45. Al-Mallah MK, Davey MR, Cocking EC. Nodulation of oilseed rape (*Brassica napus*) by *Rhizobia*. *J Exp Bot.* 1990;41:1567–72. [View ArticleGoogle Scholar](#)
46. Boddey RM, Urquiaga S, Alves BJR, Reis V. Endophytic nitrogen fixation in sugarcane: present knowledge and future applications. *Plant Soil.* 2003;252(1):139–49. [View ArticleGoogle Scholar](#)
47. Cocking EC. Endophytic colonisation of plant roots by nitrogen-fixing bacteria. *Plant Soil.* 2003;252(1):169–75. [View ArticleGoogle Scholar](#)
48. Brewin NJ. Legume root nodule symbiosis. *Biochemistry.* 2013;35(4):14–8. [Google Scholar](#)
49. Cocking EC. The challenge of establishing symbiotic nitrogen fixation in cereals. In: Emerich DW, Krishnan HW (eds) *Nitrogen fixation in crop production.* Agron Mono. 2009;52:35–64.
50. Boddey RM, Silva G, Reis VM, Alves BJR, Urquiaga S. Assessment of bacterial nitrogen fixation in grass species. In: Triplet E, editor. *Prokaryotic nitrogen fixation: a model system for analysis of a biological process.* Wymondham: Horizon Scientific; 2000. p. 705–26. [Google Scholar](#)
51. Cavalcante VA, Döbereiner J. A new acid-tolerant nitrogen-fixing bacterium associated with sugarcane. *Plant Soil.* 1988;108:23–31. [View ArticleGoogle Scholar](#)
52. Fuentes-Ramirez LE, Jimenez-Salgado T, Abarca-Ocampo IR, Caballero-Mellado J. *Acetobacter diazotrophicus*, an indoleacetic acid producing bacterium isolated from sugarcane cultivars of México. *Plant Soil.* 1993;154(2):145–50. [View ArticleGoogle Scholar](#)
53. Adriano-Anayal M, Salvador-Figueroa M, Ocampo JA, García-Romera I. Plant cell-wall degrading hydrolytic enzymes of *Gluconacetobacter diazotrophicus*. *Symbiosis.* 2005;40:151–6. [Google Scholar](#)
54. Khamar HJ, Breathwaite EK, Prasse CE, Fraley ER, Secor CR, Chibane FL, et al. Multiple roles of soluble sugars in the establishment of *Gunnera-Nostoc* endosymbiosis. *Plant Physiol.* 2010;154:1381–9. [PubMedPubMed CentralView ArticleGoogle Scholar](#)
55. Döbereiner J. History and new perspectives of diazotrophs in association with non-leguminous plants. *Symbiosis.* 1992;13:1–13. [Google Scholar](#)
56. Gillis M, Kersters K, Hoste B, Janssens D, Kroppenstedt RM, Stephan MP, et al. *Acetobacter diazotrophicus*, a nitrogen-fixing acetic acid bacterium associated with sugarcane. *Int J Syst Bacteriol.* 1989;48:327. [Google Scholar](#)
57. Sevilla M, Kennedy C. Genetic analysis of nitrogen fixation and plant-growth stimulating properties of *Acetobacter diazotrophicus*, an endophyte of sugarcane. In: Triplett EW, editor. *Prokaryotic nitrogen fixation: a model system for analysis of a biological process.* Wymondham: Horizon Scientific Press; 2000. p. 737–60. [Google Scholar](#)
58. Dong Z, Zelmer CD, Canny MJ, McCully ME, Luit B, Pan B, et al. Evidence for protection of nitrogenase from O₂ by colony structure in the aerobic diazotroph *Gluconacetobacter diazotrophicus*. *Microbiology.* 2002;148(8):2293–8. [PubMedView ArticleGoogle Scholar](#)
59. Ureta A, Nordlund S. Evidence for conformational protection of nitrogenase against oxygen in *Gluconacetobacter diazotrophicus* by a putative FeSII protein. *J Bacteriol.* 2002;148:5805–9. [View ArticleGoogle Scholar](#)

60. Cojho EH, Reis VM, Schenberg ACG, Döbereiner J. Interactions of *Acetobacter diazotrophicus* with an amylolytic yeast in nitrogen-free batch culture. *FEMS Microbiol Lett.* 1993;106:341–6. [View ArticleGoogle Scholar](#)
61. McFarlane HE, Döring A, Persson S. The cell biology of cellulose synthesis. *Annu Rev Plant Biol.* 2014;65:69–94. [PubMedView ArticleGoogle Scholar](#)
62. Postgate J. The Leeuwenhoek Lecture 1992: Bacterial Evolution and the nitrogen-fixing plant. *Philos Trans R Soc.* 1992;338(1286):409–16. [View ArticleGoogle Scholar](#)
63. Soltis DE, Soltis PS, Morgan DR, Swensen SM, Mullin BC, Dowd JM, et al. Chloroplast gene sequence data suggest a single origin of the predisposition for symbiotic fixation in angiosperms. *Proc Natl Acad Sci USA.* 1995;92:2647–51. [PubMedPubMed CentralView ArticleGoogle Scholar](#)
64. Gijsbert DA, Cornwell WK, Sprent JI, Kattge J, Kiers ET. A single evolutionary innovation drives the deep evolution of symbiotic N₂-fixation in angiosperms. *Nat Commun.* 2014;5:4087. doi: [10.1038/ncomms5087](#). [Google Scholar](#)
65. Evans JR. Nitrogen and photosynthesis in the flag leaf of wheat. *Plant Physiol.* 1983;72:297–302. [PubMedPubMed CentralView ArticleGoogle Scholar](#)
66. Tucker M. Primary nutrients and plant growth. In: SCRIBD, editor. *Essential plant nutrients.* NC, USA. North Carolina Department of Agriculture; 2004. p. 1–9.
67. Daughtry CST, Walthall CL, Kim MS, Brown de Colstoun E, McMurtrey JE III. Estimating corn leaf chlorophyll concentration from leaf and canopy reflectance. *Remote Sens Environ.* 2013;74(2):229–39. [View ArticleGoogle Scholar](#)
68. Bejandi TK, Sharifii RS, Sedghi M, Namvar A. Effects of plant density, *Rhizobium* inoculation and microelements on nodulation, chlorophyll content and yield of chickpea. *Ann Biol Res.* 2012;3(2):951–8. [Google Scholar](#)
69. Nyoki D, Ndakidemi PA. Effects of phosphorus and *Bradyrhizobium japonicum* on growth and chlorophyll content of cowpea. *Am J Exp Agric.* 2014;4:1120–36. [View ArticleGoogle Scholar](#)
70. Bambara S, Ndakidemi PA. Effects of *Rhizobium* inoculation, lime and molybdenum on photosynthesis and chlorophyll content of *Phaseolus vulgaris*. *Afr J Microbiol Res.* 2009;3(11):791–8. [Google Scholar](#)
71. Meenakshisundara M, Santhaguru K. Studies on association of arbuscular mycorrhizal fungi with *Gluconacetobacter diazotrophicus* and its effect on improvement of *Sorghum bicolor*. *Int J Curr Sci Res.* 2011;1(2):23–30. [Google Scholar](#)
72. Madhaiyan M, Saravanan VS, Jovi DB, Lee H, Thenmozhi R, Hari K, et al. Occurrence of *Gluconacetobacter diazotrophicus* in tropical and subtropical plants of Western Ghats, India. *Microbiol Res.* 2004;159(3):233–43. [PubMedView ArticleGoogle Scholar](#)
73. Momose A, Hiyama T, Nishimura K, Ishizaki N, Ishikawa S, Yamamoto M, et al. Characteristics of nitrogen fixation and nitrogen release from diazotrophic endophytes isolated from sugarcane stems. *Bull Fac Agric Niigata Univ.* 2013;66(1):1–9. [Google Scholar](#)
74. Eskin N. Colonisation of *Zea mays* by the nitrogen fixing bacterium *Gluconacetobacter diazotrophicus*. M.Sc. Thesis. The School of Graduate and Postdoctoral Studies, Western University, London, ON, Canada; 2012.
75. Sevilla M, Burris RH, Gunapala N, Kennedy C. Comparison of benefit to sugarcane plant growth and ¹⁵N₂ incorporation following inoculation of sterile plants with *Acetobacter diazotrophicus* wild-type and *Nif*-mutant strains. *Mol Plant Microbe Interact.* 2001;14(3):358–66. [PubMedView ArticleGoogle Scholar](#)
76. Tairo EV, Ndakidemi PA. *Bradyrhizobium japonicum* inoculation and phosphorus supplementation on growth and chlorophyll accumulation in Soybean. *Am J Plant Sci.* 2013;4:2281. [View ArticleGoogle Scholar](#)
77. Malik MA, Cheema MA, Khan HZ, Wahid MA. Growth and yield response of soybean to seed inoculation and varying phosphorus levels. *J Agric Res.* 2006;44:47–53. [Google Scholar](#)
78. Huang HC, Erickson RS. Effect of seed treatment with *Rhizobium leguminosarum* on *Pythium* damping-off, seedling height, root nodulation, root biomass, shoot biomass, and seed yield of pea and lentil. *J Phytopathol.* 2007;155(1):31–7. [View ArticleGoogle Scholar](#)

79. Mfilinge A, Mtei K, Ndakidemi P. Effect of Rhizobium inoculation and supplementation with phosphorus and potassium on growth and total leaf chlorophyll content of Bush Bean. *Agric Sci.* 2004;5:1413–26. [Google Scholar](#)
80. Rudresh DL, Shivaprakash MK, Prasad RD. Effect of combined application of Rhizobium, phosphate solubilizing bacterium and Trichoderma spp. on growth, nutrient uptake and yield of chickpea. *Appl Soil Ecol.* 2005;28:139–46. [View ArticleGoogle Scholar](#)
81. Ravikumar R. Growth effects of Rhizobium inoculation in some Legume plants. *Int J Curr Sci.* 2012;1:1–6. [Google Scholar](#)
82. Luna MF, Aprea J, Crespo JM, Boiard JS. Colonisation and yield promotion of tomato by *Gluconacetobacter diazotrophicus*. *Appl Soil Ecol.* 2012;61:225–9. [View ArticleGoogle Scholar](#)
83. Abudureheman A. Improving sugar beet productivity by inoculation with *Gluconacetobacter* spp. M.Sc. Thesis, Saint Mary's University, Halifax, Nova Scotia. 2012.
84. Luna MF, Galar ML, Aprea J, Molinari ML, Bioardi JL. Colonisation of sorghum and wheat by seed inoculation with *Gluconacetobacter diazotrophicus*. *Biotechnol Lett.* 2010;32:1071–6. [PubMedView ArticleGoogle Scholar](#)
85. Adriano-Anayal M, Salvador-Figueroa M, Ocampo JA, García-Romera I. Hydrolytic enzyme activities in maize and sorghum roots inoculated with *Gluconacetobacter diazotrophicus* and *Glomus intraradices*. *Soil Biol Biochem.* 2006;38:879–86. [View ArticleGoogle Scholar](#)
86. Riggs PJ, Chelius MK, Iniguez A, Kaeppler SM, Triplett EW. Enhanced maize productivity by inoculation with diazotrophic bacteria. *Aust J Plant Physiol.* 2001;28(9):829–36. [Google Scholar](#)
87. Kaschuk G, Giller K, Hungria M, Kuyper TW. Differences in photosynthetic behaviour and leaf senescence of soybean dependent on N₂ fixation or nitrate supply. *Plant Biol.* 2010;12:60–9. [PubMedView ArticleGoogle Scholar](#)
88. Bethlenfalvagy J, Abu-Shakras S, Phillips DA. Interdependence of nitrogen nutrition and photosynthesis in *Pisum sativum*: host plant response to nitrogen fixation by Rhizobium strains. *Plant Physiol.* 1978;62:131–3. [View ArticleGoogle Scholar](#)
89. Chi F, Shen S-H, Cheng H-P, Jing Y-X, Yanni YJ, Dazzo FB. Ascending migration of endophytic Rhizobia, from roots to leaves, inside rice plants and assessment of benefits to rice growth physiology. *Appl Environ Microbiol.* 2005;71(11):7271–8. [PubMedPubMed CentralView ArticleGoogle Scholar](#)
90. Rangel de Souza ALS, De Souza SA, De Oliveira MVV, Ferraz TM, Figueiredo FAMMA, Da Silva ND, et al. Endophytic colonisation of *Arabidopsis thaliana* by *Gluconacetobacter diazotrophicus* and its effect on plant growth promotion, plant physiology and activation of plant defense. *Plant Soil.* 2015;399(1):257–70. [Google Scholar](#)
91. Lynas M. *The God Species: how the planet can survive the age of humans*. London: Fourth Estate; 2011. p. 109. [Google Scholar](#)

Copyright

© The Author(s) 2017